

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
<p>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</p>					
1. REPORT DATE (DD-MM-YYYY) 17 September 2015		2. REPORT TYPE Briefing Charts		3. DATES COVERED (From - To) 24 August 2015 – 17 September 2015	
4. TITLE AND SUBTITLE Octree bin-to-bin fractional-NTC collisions				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) R. S. Martin				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER Q1AM	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/RQRS 1 Ara Drive Edwards AFB, CA 93524-7013				8. PERFORMING ORGANIZATION REPORT NO.	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/RQR 5 Pollux Drive Edwards AFB, CA 93524-7048				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S) AFRL-RQ-ED-VG-2015-350	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES For presentation at DSMC15; Kapaa, Kauai, Hawaii; 13-17 Sep 2015 PA Case Number: #15557; Clearance Date: 9/15/2015					
14. ABSTRACT Viewgraphs/Briefing Charts					
15. SUBJECT TERMS N/A					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 55	19a. NAME OF RESPONSIBLE PERSON J. Koo
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NO (include area code) N/A

OCTREE BIN-TO-BIN FRACTIONAL-NTC COLLISIONS

Robert Martin

ERC Inc.,
SPACECRAFT PROPULSION BRANCH
AIR FORCE RESEARCH LABORATORY
EDWARDS AIR FORCE BASE, CA USA

DSMC 2015

Distribution A: Approved for public release;
Distribution Unlimited;
AFTC/PA clearance No. TBD



U.S. AIR FORCE





- 1 BACKGROUND
- 2 FRACTIONAL COLLISIONS
- 3 BIN-TO-BIN FRACTIONAL-NTC
- 4 CONCLUSION



IMPORTANCE OF COLLISION PHYSICS


Important Collisions in Spacecraft Propulsion:

- Discharge and Breakdown in FRC
- Collisional Radiative Cooling/Ionization
- Combustion Chemistry


Common Features in Spacecraft Collisions:

- Relevant Densities Spanning Many Orders of Magnitude — 6+
- Transitions from Collisional to Collisionless
- Tiny Early e^- or Radical Populations Critical to Induction Delay
- Many types of Inelastic Collisions with Unknown Effects on Distribution Shapes

Shock Ionization



Kapper & Cambier, J. Appl. Phys. 109, (2011)





IMPORTANCE OF COLLISION PHYSICS

Important Collisions in Spacecraft Propulsion:


- Discharge and Breakdown in FRC
- Collisional Radiative Cooling/Ionization
- Combustion Chemistry

Common Features in Spacecraft Collisions:


- Relevant Densities Spanning Many Orders of Magnitude — 6+
- Transitions from Collisional to Collisionless
- Tiny Early e^- or Radical Populations Critical to Induction Delay
- Many types of Inelastic Collisions with Unknown Effects on Distribution Shapes

Need Low Noise & High Dynamic Range
Collision Algorithms

Shock Ionization



Kapper & Cambier, J. Appl. Phys. 109, (2011)





Previous Collision Methods:

- Monte Carlo Collisions (MCC)
 - Particles Collide with Background “Fluid”
 - Often Used in Plasma/PIC Simulation
 - Ion- e^- Collisions Assume Stationary Ions
 - No Conservation/Detailed Balance
- Direct Simulation Monte Carlo Collisions (DSMC)
 - Most Modern Versions use No-Time Counter (NTC) Method
 - Conservative/Reversible Collision
 - Satisfies Detailed Balance
 - Subset of Possible Collisions Sampled
 - Random Selection vs Z_{ij} for All/Nothing Collision

All Random Flip vs Number of Collisions: $Z_{ij} = \frac{n_i n_j}{2} \langle \sigma v \rangle dt$



VARIABLE WEIGHTS FOR DYNAMIC RANGE

Continuum to Discrete Representation:

- Many Particles \rightsquigarrow Continuous Distribution





VARIABLE WEIGHTS FOR DYNAMIC RANGE

Continuum to Discrete Representation:

- Many Particles \rightsquigarrow Continuous Distribution
- Discretized VDF Yields Vlasov
But Collision Integral Still a Problem





Continuum to Discrete Representation:

- Many Particles \rightsquigarrow Continuous Distribution
- Discretized VDF Yields Vlasov
But Collision Integral Still a Problem
- Particle Methods VDF to Delta Function Set
- Collisions between Discrete Velocities






VARIABLE WEIGHTS FOR DYNAMIC RANGE

Continuum to Discrete Representation:

- Many Particles \rightsquigarrow Continuous Distribution
- Discretized VDF Yields Vlasov
But Collision Integral Still a Problem
- Particle Methods VDF to Delta Function Set
- Collisions between Discrete Velocities
- But Poorly Resolved Tail
(Tail Critical to Inelastic Collisions)





VARIABLE WEIGHTS FOR DYNAMIC RANGE

Continuum to Discrete Representation:

- Many Particles \rightsquigarrow Continuous Distribution
- Discretized VDF Yields Vlasov
But Collision Integral Still a Problem
- Particle Methods VDF to Delta Function Set
- Collisions between Discrete Velocities
- But Poorly Resolved Tail
(Tail Critical to Inelastic Collisions)
- Variable Weights Permit Extra DOF in Tails






VARIABLE WEIGHTS FOR DYNAMIC RANGE

Continuum to Discrete Representation:

- Many Particles \rightsquigarrow Continuous Distribution
- Discretized VDF Yields Vlasov
But Collision Integral Still a Problem
- Particle Methods VDF to Delta Function Set
- Collisions between Discrete Velocities
- But Poorly Resolved Tail
(Tail Critical to Inelastic Collisions)
- Variable Weights Permit Extra DOF in Tails

Variable Weight “All-or-Nothing” Collisions?






VARIABLE WEIGHTS FOR DYNAMIC RANGE

Continuum to Discrete Representation:

- Many Particles \rightsquigarrow Continuous Distribution
- Discretized VDF Yields Vlasov
But Collision Integral Still a Problem
- Particle Methods VDF to Delta Function Set
- Collisions between Discrete Velocities
- But Poorly Resolved Tail
(Tail Critical to Inelastic Collisions)
- Variable Weights Permit Extra DOF in Tails

Variable Weight “All-or-Nothing” Collisions?





VARIABLE WEIGHTS FOR DYNAMIC RANGE


Continuum to Discrete Representation:

- Many Particles \rightsquigarrow Continuous Distribution
- Discretized VDF Yields Vlasov
But Collision Integral Still a Problem
- Particle Methods VDF to Delta Function Set
- Collisions between Discrete Velocities
- But Poorly Resolved Tail
(Tail Critical to Inelastic Collisions)
- Variable Weights Permit Extra DOF in Tails

Variable Weight “All-or-Nothing” Collisions?

Physically Inconsistent!

(Mixing Violates Momentum/Energy Conservation)





REVIEW OF FRACTIONAL COLLISIONS

NTC Collisions:

- (Collision Rate Volume):(Cell Volume)

Fractional-NTC Collisions:

$$Z_{ij} = \frac{n_i n_j}{2} \langle \sigma v \rangle_{ij} dt = \frac{w_i w_j}{2V_{cell}^2} \langle \sigma v \rangle_{ij} dt$$



REVIEW OF FRACTIONAL COLLISIONS

NTC Collisions:

- (Collision Rate Volume):(Cell Volume)
- Select Fraction of $\frac{1}{2}N^2$ Possible

Fractional-NTC Collisions:

$$Z_{ij} = \frac{n_i n_j}{2} \langle \sigma v \rangle_{ij} dt = \frac{w_i w_j}{2V_{cell}^2} \langle \sigma v \rangle_{ij} dt$$

$$P_{ij} = w \langle \sigma v \rangle_{ij} dt / V_{cell}$$

$$P_{max} = w \langle \sigma v \rangle_{ij}^{max} dt / V_{cell}$$

$$N_{select} = \frac{N_p^2}{2} F_n \langle \sigma v \rangle_{ij}^{max} dt / V_{cell}$$



REVIEW OF FRACTIONAL COLLISIONS

NTC Collisions:

- (Collision Rate Volume):(Cell Volume)
- Select Fraction of $\frac{1}{2}N^2$ Possible
- Probability of Event Ratio of Volumes

Fractional-NTC Collisions:

$$Z_{ij} = \frac{n_i n_j}{2} \langle \sigma v \rangle_{ij} dt = \frac{w_i w_j}{2V_{cell}^2} \langle \sigma v \rangle_{ij} dt$$

$$P_{ij} = w \langle \sigma v \rangle_{ij} dt / V_{cell}$$

$$P_{max} = w \langle \sigma v \rangle_{ij}^{max} dt / V_{cell}$$

$$N_{select} = \frac{N_p^2}{2} F_n \langle \sigma v \rangle_{ij}^{max} dt / V_{cell}$$

Collide if:

$$\text{Rand}(1) < \frac{N_{collide}}{N_{select}} = \frac{P_{ij}}{P_{max}} = \frac{\langle \sigma v \rangle_{ij}}{\langle \sigma v \rangle_{ij}^{max}}$$



REVIEW OF FRACTIONAL COLLISIONS

NTC Collisions:

- (Collision Rate Volume):(Cell Volume)
- Select Fraction of $\frac{1}{2}N^2$ Possible
- Probability of Event Ratio of Volumes
- Correct Non-Equilibrium Frequency

Fractional-NTC Collisions:

$$Z_{ij} = \frac{n_i n_j}{2} \langle \sigma v \rangle_{ij} dt = \frac{w_i w_j}{2V_{cell}^2} \langle \sigma v \rangle_{ij} dt$$

$$P_{ij} = w \langle \sigma v \rangle_{ij} dt / V_{cell}$$

$$P_{max} = w \langle \sigma v \rangle_{ij}^{max} dt / V_{cell}$$

$$N_{select} = \frac{N_p^2}{2} F_n \langle \sigma v \rangle_{ij}^{max} dt / V_{cell}$$

Collide if:

$$\text{Rand}(1) < \frac{N_{collide}}{N_{select}} = \frac{P_{ij}}{P_{max}} = \frac{\langle \sigma v \rangle_{ij}}{\langle \sigma v \rangle_{ij}^{max}}$$



REVIEW OF FRACTIONAL COLLISIONS

NTC Collisions:

- (Collision Rate Volume):(Cell Volume)
- Select Fraction of $\frac{1}{2}N^2$ Possible
- Probability of Event Ratio of Volumes
- Correct Non-Equilibrium Frequency

Fractional-NTC Collisions:

- Select f by Cost/Accuracy Tradeoff

$$Z_{ij} = \frac{n_i n_j}{2} \langle \sigma v \rangle_{ij} dt = \frac{w_i w_j}{2V_{cell}^2} \langle \sigma v \rangle_{ij} dt$$

$$P_{ij} = w \langle \sigma v \rangle_{ij} dt / V_{cell}$$

$$P_{max} = w \langle \sigma v \rangle_{ij}^{max} dt / V_{cell}$$

$$N_{select} = \frac{N_p^2}{2} F_n \langle \sigma v \rangle_{ij}^{max} dt / V_{cell}$$

Collide if:

$$\text{Rand}(1) < \frac{N_{collide}}{N_{select}} = \frac{P_{ij}}{P_{max}} = \frac{\langle \sigma v \rangle_{ij}}{\langle \sigma v \rangle_{ij}^{max}}$$

$$N_{select} = f N_p$$



REVIEW OF FRACTIONAL COLLISIONS

NTC Collisions:

- (Collision Rate Volume):(Cell Volume)
- Select Fraction of $\frac{1}{2}N^2$ Possible
- Probability of Event Ratio of Volumes
- Correct Non-Equilibrium Frequency

Fractional-NTC Collisions:

- Select f by Cost/Accuracy Tradeoff
- Collision Δw Scaled for Skipped

$$Z_{ij} = \frac{n_i n_j}{2} \langle \sigma v \rangle_{ij} dt = \frac{w_i w_j}{2V_{cell}^2} \langle \sigma v \rangle_{ij} dt$$

$$P_{ij} = w \langle \sigma v \rangle_{ij} dt / V_{cell}$$

$$P_{max} = w \langle \sigma v \rangle_{ij}^{max} dt / V_{cell}$$

$$N_{select} = \frac{N_p^2}{2} F_n \langle \sigma v \rangle_{ij}^{max} dt / V_{cell}$$

Collide if:

$$\text{Rand}(1) < \frac{N_{collide}}{N_{select}} = \frac{P_{ij}}{P_{max}} = \frac{\langle \sigma v \rangle_{ij}}{\langle \sigma v \rangle_{ij}^{max}}$$

$$N_{select} = f N_p$$

$$\Delta w_{ij} = \frac{N_p^2 / 2}{N_{select}} Z_{ij}$$



REVIEW OF FRACTIONAL COLLISIONS

NTC Collisions:

- (Collision Rate Volume):(Cell Volume)
- Select Fraction of $\frac{1}{2}N^2$ Possible
- Probability of Event Ratio of Volumes
- Correct Non-Equilibrium Frequency

Fractional-NTC Collisions:

- Select f by Cost/Accuracy Tradeoff
- Collision Δw Scaled for Skipped
- Add Particles & Original Reduced

$$Z_{ij} = \frac{n_i n_j}{2} \langle \sigma v \rangle_{ij} dt = \frac{w_i w_j}{2V_{cell}^2} \langle \sigma v \rangle_{ij} dt$$

$$P_{ij} = w \langle \sigma v \rangle_{ij} dt / V_{cell}$$

$$P_{max} = w \langle \sigma v \rangle_{ij}^{max} dt / V_{cell}$$

$$N_{select} = \frac{N_p^2}{2} F_n \langle \sigma v \rangle_{ij}^{max} dt / V_{cell}$$

Collide if:

$$\text{Rand}(1) < \frac{N_{collide}}{N_{select}} = \frac{P_{ij}}{P_{max}} = \frac{\langle \sigma v \rangle_{ij}}{\langle \sigma v \rangle_{ij}^{max}}$$

$$N_{select} = f N_p$$

$$\Delta w_{ij} = \frac{N_p^2 / 2}{N_{select}} Z_{ij}$$

$$w_i = w_i - \Delta w_{ij} \ \& \ w_j = w_j - \Delta w_{ij}$$

$$w_{(N_p+1)} = \Delta w_{ij} \ \& \ w_{(N_p+2)} = \Delta w_{ij}$$



REVIEW OF FRACTIONAL COLLISIONS


NTC Collisions:

- (Collision Rate Volume):(Cell Volume)
- Select Fraction of $\frac{1}{2}N^2$ Possible
- Probability of Event Ratio of Volumes
- Correct Non-Equilibrium Frequency

Fractional-NTC Collisions:

- Select f by Cost/Accuracy Tradeoff
- Collision Δw Scaled for Skipped
- Add Particles & Original Reduced
- **+2 Particles/Collision! → Must Merge**

$$Z_{ij} = \frac{n_i n_j}{2} \langle \sigma v \rangle_{ij} dt = \frac{w_i w_j}{2V_{cell}^2} \langle \sigma v \rangle_{ij} dt$$



$$w_i = w_i - \Delta w_{ij} \ \& \ w_j = w_j - \Delta w_{ij}$$


$$w_{(N_p+1)} = \Delta w_{ij} \ \& \ w_{(N_p+2)} = \Delta w_{ij}$$



REVIEW OF CONSERVATIVE MERGE

Merge to Pair → DOF for Conservation:

- $(n+2):2$ yields Exact Mass, Momentum, and Kinetic Energy Conservation
- Applied Spatially also Shown to Conserve Electrostatic Energy
- Though Energy Conserving, Still Thermalizes VDF



$$w_{cell} = \sum_i^{(n+2)} w_i$$

$$\vec{v} = \frac{1}{w_{cell}} \sum_i^{(n+2)} w_i \vec{v}_i$$

$$\overline{V^2} = \frac{1}{w_{cell}} \sum_i^{(n+2)} w_i \left(\vec{v}_i - \vec{v} \right)^2$$

$$w_{(a/b)} = w_m/2$$

$$\vec{v}_{(a/b)} = \vec{v} \pm \hat{\mathcal{R}} \sqrt{\overline{V^2}}$$

$$\text{Similarly: } \vec{x}_{(a/b)} = \vec{x} \pm \hat{\mathcal{R}} \sqrt{\overline{X^2}}$$




REVIEW OF CONSERVATIVE MERGE

Merge to Pair → DOF for Conservation:

- $(n+2):2$ yields Exact Mass, Momentum, and Kinetic Energy Conservation
- Applied Spatially also Shown to Conserve Electrostatic Energy
- Though Energy Conserving, Still Thermalizes VDF

Selection of Near Neighbors in VDF Limits Thermalization

(\approx Near Neighbor Pairs in 2:1 Merges that Limit Numerical Cooling)



$$w_{cell} = \sum_i^{(n+2)} w_i$$

$$\vec{v} = \frac{1}{w_{cell}} \sum_i^{(n+2)} w_i \vec{v}_i$$

$$\overline{V^2} = \frac{1}{w_{cell}} \sum_i^{(n+2)} w_i \left(\vec{v}_i - \vec{v} \right)^2$$

$$w_{(a/b)} = w_m/2$$

$$\vec{v}_{(a/b)} = \vec{v} \pm \hat{\mathcal{R}} \sqrt{\overline{V^2}}$$

$$\text{Similarly: } \vec{x}_{(a/b)} = \vec{x} \pm \hat{\mathcal{R}} \sqrt{\overline{X^2}}$$



REVIEW OF CONSERVATIVE MERGE


Merge to Pair \rightarrow DOF for Conservation:

- $(n+2):2$ yields Exact Mass, Momentum, and Kinetic Energy Conservation
- Applied Spatially also Shown to Conserve Electrostatic Energy
- Though Energy Conserving, Still Thermalizes VDF

Selection of Near Neighbors in VDF Limits Thermalization

(\approx Near Neighbor Pairs in 2:1 Merges that Limit Numerical Cooling)

Octree Velocity Bins




Efficient Neighbor Selection



0D-THERMALIZATION

Bi-Maxwellian Thermalization Results




Comparison of 10x Runs from Same Initial Distribution




0D-THERMALIZATION

Bi-Maxwellian Thermalization Results

Original-NTC



Fractional-NTC




Mean and RMS Fluctuation of Sample Runs

Fluctuations Level Tuneable with f Independent of Particles Count



0D-THERMALIZATION

Bi-Maxwellian Thermalization Results




Fluctuations Level Tuneable with f Independent of Particles Count



- Initial Bi-Maxwellian Distribution in Potential Well




- Initial Bi-Maxwellian Distribution in Potential Well
- NTC Collisions Results in Beam Thermalization





COLLISIONAL BEAMS IN POTENTIAL WELL


- Initial Bi-Maxwellian Distribution in Potential Well
- NTC Collisions Results in Beam Thermalization
- Fractional-NTC Collisions Produce Same Behavior





COLLISIONAL BEAMS IN POTENTIAL WELL


- Initial Bi-Maxwellian Distribution in Potential Well
- NTC Collisions Results in Beam Thermalization
- Fractional-NTC Collisions Produce Same Behavior
- Particles/Cell Dramatically Different





COLLISIONAL BEAMS IN POTENTIAL WELL


- Initial Bi-Maxwellian Distribution in Potential Well
- NTC Collisions Results in Beam Thermalization
- Fractional-NTC Collisions Produce Same Behavior
- Particles/Cell Dramatically Different
- Fringe Extends to Lower Densities with Variable Weights






COLLISIONAL BEAMS IN POTENTIAL WELL

- Initial Bi-Maxwellian Distribution in Potential Well
- NTC Collisions Results in Beam Thermalization
- Fractional-NTC Collisions Produce Same Behavior
- Particles/Cell Dramatically Different
- Fringe Extends to Lower Densities with Variable Weights
- Relative 'Error' Unknown without Analytical Solution or High Fidelity Simulation






- Larger $N_{select} \rightarrow$ Better Approx. of Collision Integral






- Larger $N_{select} \rightarrow$ Better Approx. of Collision Integral
- f-NTC Produces 2x-Particles per $N_{select} = f N_p$
- Particle Memory Requires $\propto N_{max} \rightarrow (1 + 2f)N_{max}$






- Larger $N_{select} \rightarrow$ Better Approx. of Collision Integral
- f-NTC Produces 2x-Particles per $N_{select} = f N_p$
- Particle Memory Requires $\propto N_{max} \rightarrow (1 + 2f)N_{max}$
- For DSMC-like Results, $f \approx O(1)$






- Larger $N_{select} \rightarrow$ Better Approx. of Collision Integral
- f-NTC Produces 2x-Particles per $N_{select} = f N_p$
- Particle Memory Requires $\propto N_{max} \rightarrow (1 + 2f)N_{max}$
- For DSMC-like Results, $f \approx O(1)$
- Time Accurate or Dense Simulations, $f \approx O(10)+?$






- Larger $N_{select} \rightarrow$ Better Approx. of Collision Integral
- f-NTC Produces 2x-Particles per $N_{select} = f N_p$
- Particle Memory Requires $\propto N_{max} \rightarrow (1 + 2f)N_{max}$
- For DSMC-like Results, $f \approx O(1)$
- Time Accurate or Dense Simulations, $f \approx O(10)+?$
- Merge Contracts back to $O(N_{max})$ Particles
- Merge Immediately after Collide per Spatial Cell?..
- Sort for Merge still $\propto (1 + 2f) \log(1 + 2f)?$





ISSUE WITH COLLIDE THEN MERGE

- Larger $N_{select} \rightarrow$ Better Approx. of Collision Integral
- f-NTC Produces 2x-Particles per $N_{select} = f N_p$
- Particle Memory Requires $\propto N_{max} \rightarrow (1 + 2f)N_{max}$
- For DSMC-like Results, $f \approx O(1)$
- Time Accurate or Dense Simulations, $f \approx O(10)+?$
- Merge Contracts back to $O(N_{max})$ Particles
- Merge Immediately after Collide per Spatial Cell?..
- Sort for Merge still $\propto (1 + 2f) \log(1 + 2f)?$
- Combine Collision and Merge in Single Step?





COLLIDE TO BINS

- Fractional Collision as Rate Equation

$$\begin{bmatrix} \vdots \\ \dot{w}_i \\ \vdots \\ \dot{w}_j \\ \vdots \\ \dot{w}_{i'} \\ \vdots \\ \dot{w}_{j'} \\ \vdots \end{bmatrix} = \sum_{k=1}^{N_{select}} \begin{bmatrix} \vdots \\ -w_i \langle \sigma v \rangle_{ij}^k w_j \\ \vdots \\ -w_i \langle \sigma v \rangle_{ij}^k w_j \\ \vdots \\ w_i \langle \sigma v \rangle_{ij}^k w_j \\ \vdots \\ w_i \langle \sigma v \rangle_{ij}^k w_j \\ \vdots \end{bmatrix}$$



COLLIDE TO BINS

- Fractional Collision as Rate Equation
- Bin Moments needed for Particle Pairs

$$\begin{bmatrix} \dot{w}_i \\ \dot{w}_j \\ \dot{w}_{i'} \\ \dot{w}_{j'} \\ - \\ (\dot{wv})_i \\ (\dot{wv})_j \\ (\dot{wv})_{i'} \\ (\dot{wv})_{j'} \\ - \\ (\dot{wv^2})_i \\ (\dot{wv^2})_j \\ (\dot{wv^2})_{i'} \\ (\dot{wv^2})_{j'} \end{bmatrix} = \sum_{k=1}^{N_{select}} \begin{bmatrix} -\Delta w_{ij} \\ -\Delta w_{ij} \\ \Delta w_{ij} \\ \Delta w_{ij} \\ - \\ -\Delta w_{ij}v_i \\ -\Delta w_{ij}v_j \\ \Delta w_{ij}v_{i'} \\ \Delta w_{ij}v_{j'} \\ - \\ -\Delta w_{ij}v_i^2 \\ -\Delta w_{ij}v_j^2 \\ \Delta w_{ij}v_{i'}^2 \\ \Delta w_{ij}v_{j'}^2 \end{bmatrix}$$



COLLIDE TO BINS

- Fractional Collision as Rate Equation
- Bin Moments needed for Particle Pairs
- Particle Pairs (i,j) Picked Randomly
- DSMC-like Collision (VHS,VSS,etc.)
Random $\chi, \theta \rightarrow (v_{i'}, v_{j'})$

$$\begin{bmatrix} \dot{w}_i \\ \dot{w}_j \\ \dot{w}_{i'} \\ \dot{w}_{j'} \\ - \\ (\dot{wv})_i \\ (\dot{wv})_j \\ (\dot{wv})_{i'} \\ (\dot{wv})_{j'} \\ - \\ (\dot{wv^2})_i \\ (\dot{wv^2})_j \\ (\dot{wv^2})_{i'} \\ (\dot{wv^2})_{j'} \end{bmatrix} = \sum_{k=1}^{N_{select}} \begin{bmatrix} -\Delta w_{ij} \\ -\Delta w_{ij} \\ \Delta w_{ij} \\ \Delta w_{ij} \\ - \\ -\Delta w_{ij}v_i \\ -\Delta w_{ij}v_j \\ \Delta w_{ij}v_{i'} \\ \Delta w_{ij}v_{j'} \\ - \\ -\Delta w_{ij}v_i^2 \\ -\Delta w_{ij}v_j^2 \\ \Delta w_{ij}v_{i'}^2 \\ \Delta w_{ij}v_{j'}^2 \end{bmatrix}$$



COLLIDE TO BINS

- Fractional Collision as Rate Equation
- Bin Moments needed for Particle Pairs
- Particle Pairs (i,j) Picked Randomly
- DSMC-like Collision (VHS,VSS,etc.)
Random $\chi, \theta \rightarrow (v_{i'}, v_{j'})$
- Octree to Find i' and j' Bins
 $8^L \rightarrow$ Few Levels to Search

$$\begin{bmatrix} \dot{w}_i \\ \dot{w}_j \\ \dot{w}_{i'} \\ \dot{w}_{j'} \\ - \\ (\dot{wv})_i \\ (\dot{wv})_j \\ (\dot{wv})_{i'} \\ (\dot{wv})_{j'} \\ - \\ (\dot{wv^2})_i \\ (\dot{wv^2})_j \\ (\dot{wv^2})_{i'} \\ (\dot{wv^2})_{j'} \end{bmatrix} = \sum_{k=1}^{N_{select}} \begin{bmatrix} -\Delta w_{ij} \\ -\Delta w_{ij} \\ \Delta w_{ij} \\ \Delta w_{ij} \\ - \\ -\Delta w_{ij}v_i \\ -\Delta w_{ij}v_j \\ \Delta w_{ij}v_{i'} \\ \Delta w_{ij}v_{j'} \\ - \\ -\Delta w_{ij}v_i^2 \\ -\Delta w_{ij}v_j^2 \\ \Delta w_{ij}v_{i'}^2 \\ \Delta w_{ij}v_{j'}^2 \end{bmatrix}$$



COLLIDE TO BINS

- Fractional Collision as Rate Equation
- Bin Moments needed for Particle Pairs
- Particle Pairs (i,j) Picked Randomly
- DSMC-like Collision (VHS,VSS,etc.)
Random $\chi, \theta \rightarrow (v_{i'}, v_{j'})$
- Octree to Find i' and j' Bins
 $8^L \rightarrow$ Few Levels to Search

Conserve Mass, Momentum, and Energy
Memory Constant Independent of N^{select}

$$\begin{bmatrix} \dot{w}_i \\ \dot{w}_j \\ \dot{w}_{i'} \\ \dot{w}_{j'} \\ - \\ (\dot{wv})_i \\ (\dot{wv})_j \\ (\dot{wv})_{i'} \\ (\dot{wv})_{j'} \\ - \\ (\dot{wv^2})_i \\ (\dot{wv^2})_j \\ (\dot{wv^2})_{i'} \\ (\dot{wv^2})_{j'} \end{bmatrix} = \sum_{k=1}^{N_{select}} \begin{bmatrix} -\Delta w_{ij} \\ -\Delta w_{ij} \\ \Delta w_{ij} \\ \Delta w_{ij} \\ - \\ -\Delta w_{ij}v_i \\ -\Delta w_{ij}v_j \\ \Delta w_{ij}v_{i'} \\ \Delta w_{ij}v_{j'} \\ - \\ -\Delta w_{ij}v_i^2 \\ -\Delta w_{ij}v_j^2 \\ \Delta w_{ij}v_{i'}^2 \\ \Delta w_{ij}v_{j'}^2 \end{bmatrix}$$



Non-linear Equation in Terms of Weights:

$$\frac{\delta Q}{\delta t} = F(Q)$$

Amenable to Time Marching Schemes?




Non-linear Equation in Terms of Weights:

$$\frac{\delta Q}{\delta t} = F(Q)$$

Explicit:

$$\delta Q = \Delta t F(Q)$$






Non-linear Equation in Terms of Weights:

$$\frac{\delta Q}{\delta t} = F(Q)$$

Predictor Corrector:

$$\overline{\delta Q} = \frac{\Delta t}{2} (F(Q^0) + F(Q^0 + \delta Q))$$

(Iterate for δQ ?)





Non-linear Equation in Terms of Weights:

$$\frac{\delta Q}{\delta t} = F(Q)$$

Linearized Crank-Nicolson:

$$\delta Q = \Delta t \left(F(Q^0) + \frac{1}{2} \left. \frac{\delta F}{\delta Q} \right|^0 \delta Q \right)$$

$$\delta Q = \Delta t \left(I - \frac{\Delta t}{2} \left. \frac{\delta F}{\delta Q} \right|^0 \right)^{-1} F(Q^0)$$



Non-linear Equation in Terms of Weights:

$$\frac{\delta Q}{\delta t} = F(Q)$$

Linearized Crank-Nicolson:

$$\delta Q = \Delta t \left(F(Q^0) + \frac{1}{2} \frac{\delta F}{\delta Q} \Big|_0 \delta Q \right)$$

$$\delta Q = \Delta t \left(I - \frac{\Delta t}{2} \frac{\delta F}{\delta Q} \Big|_0 \right)^{-1} F(Q^0)$$

But Complex in v_i, v_j in Terms of $Q...$



Non-linear Equation in Terms of Weights:

$$\frac{\delta Q}{\delta t} = F(Q)$$

Linearized Crank-Nicholson:


$$\delta Q^{(w)} = \Delta t \left(F(Q^{(w),0}) + \frac{1}{2} \frac{\delta F^{(w)}}{\delta Q^{(w)}} \bigg|_0 \delta Q^{(w)} \right)$$

$$\delta Q^{(w)} = \Delta t \left(I - \frac{\Delta t}{2} \frac{\delta F^{(w)}}{\delta Q^{(w)}} \bigg|_0 \right)^{-1} F(Q^{(w),0})$$

$$\delta Q^{(M)} \approx \Delta t \left(F(Q^{(M),0}) + \frac{1}{2} \frac{\delta F^{(M)}}{\delta Q^{(w)}} \bigg|_0 \delta Q^{(w)} \right)$$

But Complex in v_i, v_j in Terms of $Q...$

First Assume Primary Dependence on $\delta w..?$





Marching Worse than Original F-NTC?

$$\frac{\delta Q}{\delta t} = F(Q)$$

Approximation of Random Discrete Jump
Process



DIRECT TIME MARCHING?

Marching Worse than Original F-NTC?

$$\frac{\delta Q}{\delta t} = F(Q)$$

Continuous:


$$\delta Q \approx \Delta t F(Q) \rightarrow \delta Q = \int_0^{\Delta t} F(Q(t)) dt$$

Order of Jumps is Random

Approximates Nature as $N^{select} \rightarrow \infty$ Increase

$f \rightarrow$ Smaller Jumps Independent of Δt

Continuous Update of $Q \approx$ VDF Evolution





- Standard Collision Incompatible with Variable Weight
- Fractional-NTC Option for Variable Weight Collision
- Bin-To-Bin F-NTC Potentially Alleviates Memory Constraints
- Bin-To-Bin also Allows Route to Advanced Time Marching
- Preliminary Advanced Time Marching Requires Additional Study
- Verification vs. Standard Shock Cases/etc. Needed



END



Thank You

Questions?